
Bovine Spongiform Encephalopathy (BSE) in Animals 2016

In 2016, one out of approx. 60,000 total tests carried out for BSE in bovines were positive. The number of BSE cases peaked in 2002 (333 cases), but since then a steady decline in incidence has been observed. This is mainly attributed to older animals in the national herd being replaced by younger animals that have not been exposed to contaminated feed. Since 2008, no BSE incidents have been detected in cattle younger than 60 months of age.



Transmissible Spongiform Encephalopathy (TSE) in Animals 2016

In 2016, 20,000 sheep and 130 goats were tested for scrapie. Of these, one flock was positive for classical scrapie and one was positive for atypical scrapie.

Sources of data:

Department of Agriculture, Food and the Marine (DAFM): Annual Report 2016. https://www.agriculture.gov.ie/media/migration/publications/2017/FinalDAFM2016AnnualReport0 https://www.agriculture.gov.ie/media/migration/publications/2017/FinalDAFM2016AnnualReport0 https://www.agriculture.gov.ie/media/migration/publications/2017/FinalDAFM2016AnnualReport0 https://www.agriculture.gov.ie/media/migration/publications/2017/FinalDAFM2016AnnualReport0 https://www.agriculture.gov.ie/media/migration/publications/2017/FinalDAFM2016AnnualReport0 https://www.agriculture.gov.ie/migration/publications/2017/FinalDAFM2016AnnualReport0 https://www.agriculture.gov.ie/migration/ <a href="