

Other Zoonoses in Animals 2016

Table 1: *Yersinia* in animals

| Animals | Tested | Positive | Species |
|-------------------------|---------------|------------------|--|
| Suspect Sampling | | | |
| Birds | 82 | 0 | |
| Cats | 1 | 0 | |
| Cattle (bovine) | 6167 | 2 (0.03%) | <i>Yersinia pseudotuberculosis</i> (n=2) |
| Deer (farmed) | 4 | 0 | |
| Dogs | 129 | 0 | |
| Goats | 33 | 0 | |
| Pigs (fattening) | 547 | 0 | |
| Poultry | 326 | 0 | |
| Rabbits | 11 | 0 | |
| Sheep | 2989 | 0 | |
| Solipeds | 53 | 0 | |
| Total | 10,342 | 2 (0.02%) | <i>Yersinia pseudotuberculosis</i> (n=2) |

Source: DAFM

Table 2: *Trichinella* in animals

| Animals | Tested | Positive |
|---------------------------|--------------|----------|
| Objective Sampling | | |
| Pigs (fattening) | 3385 | 0 |
| Solipeds | 140 | 0 |
| Total | 3,525 | 0 |

Source: DAFM

Table 3. *Coxiella* in animals

| Animals | Tested | Positive (%) | Species |
|-------------------------|------------|------------------|---------------------------------|
| Suspect Sampling | | | |
| Cattle (bovine) | 97 | 11 (11.3%) | <i>Coxiella burnetii</i> (n=11) |
| Sheep | 475 | 5 (1.1%) | <i>Coxiella burnetii</i> (n=5) |
| Total | 572 | 16 (2.8%) | <i>Coxiella burnetii</i> (n=16) |

Source: DAFM

Source of data:

Department of Agriculture, Food and the Marine (DAFM)