



Salt and Health: Review of the Scientific Evidence and Recommendations for Public Policy in Ireland





Salt and Health: Review of the Scientific Evidence and Recommendations for Public Policy in Ireland



**Published by:
Food Safety Authority of Ireland
Abbey Court
Lower Abbey Street
Dublin 1**

**Telephone: +353 1 817 1300
Facsimile: +353 1 817 1301
E-mail: info@fsai.ie
Website: www.fsai.ie**

©2005



I.S. EN ISO 9001:2000

ISBN 1-904465-23-4

CONTENTS

| | |
|---|-----------|
| FOREWORD | 1 |
| Members of the Scientific Committee | 1 |
| Members of the Nutrition Sub-committee | 1 |
| SUMMARY | 2 |
| RECOMMENDATIONS | 4 |
| General Recommendations | 4 |
| Recommendations for the FSAI | 4 |
| Recommendations for Other Agencies | 5 |
| Recommendations for the Food Industry | 5 |
| BACKGROUND TO THIS REPORT | 6 |
| 1. HYPERTENSION AND CARDIOVASCULAR DISEASE IN IRELAND | 7 |
| Definition of Hypertension | 7 |
| Prevalence of Hypertension | 7 |
| Hypertension and Cardiovascular Disease | 8 |
| 2. OVERVIEW OF EVIDENCE ON SALT AND BLOOD PRESSURE | 9 |
| Meta-analyses of Salt Restriction Intervention Studies | 10 |
| Salt Restriction in the Elderly | 11 |
| The Dash Sodium Trial | 11 |
| Salt, Left Ventricular Hypertrophy (Cardiac Enlargement) and Stroke | 12 |
| 3. THE ROLE OF SALT RELATIVE TO OTHER FACTORS IN THE DEVELOPMENT OF HYPERTENSION | 13 |
| Causes of Hypertension | 13 |
| Salt Taste Thresholds | 15 |



| | |
|--|-----------|
| 4. RECOMMENDED LEVELS OF SODIUM INTAKE | 16 |
| Potential Interventions to Reduce Population Salt Intake | 16 |
| 5. CONCLUSIONS | 18 |
| 6. MAJOR RECOMMENDATIONS | 19 |
| General Recommendations | 19 |
| Recommendations for the FSAI | 19 |
| Recommendations for Other Agencies | 20 |
| Recommendations for the Food Industry | 20 |
| REFERENCES | 21 |



FOREWORD

This report was prepared by the Nutrition Sub-committee and adopted by the Scientific Committee for presentation to the Food Safety Authority of Ireland (FSAI). It aims to provide the FSAI and other stakeholders with an overview of the science and related issues surrounding the problem of excessive salt in the diet. It also provides recommendations for all stakeholders to address and resolve this public health issue.

Members of the Scientific Committee (up to Nov 2004)

Prof. Albert Flynn (Chair)
University College, Cork

Prof. Martin Cormican
National University of Ireland, Galway

Ms Mary Falvey
Southern Health Board

Dr Clíodhna Foley-Nolan
Southern Health Board

Dr Michael Gunn
Department of Agriculture and Food

Prof. Colin Hill
University College, Cork

Dr Mark Lynch
Department of Agriculture and Food

Ms Siobhan McEvoy
Department of Health and Children

Prof. Brian McKenna
University College, Dublin

Dr Marlene Proctor
DIT, Cathal Brugha Street

Prof. Jim Roche
University College, Dublin

Prof. Michael P Ryan
University College, Dublin

Dr James Sheridan
Teagasc

Prof. Peter Smith
National University of Ireland, Galway

Dr Maire Walsh
State Laboratory

Members of the Nutrition Sub-committee (up to Nov 2004)

Prof. Albert Flynn (Chair)
University College Cork, Cork

Dr Sharon Friel
National University of Ireland, Galway

Dr John Kearney
Trinity College, Dublin

Dr Mairead Kiely
University College, Cork

Dr Helene McNulty
University of Ulster, Coleraine

Ms Ursula O'Dwyer
Department of Health and Children

Prof. Ivan Perry
University College, Cork

Dr Helen Roche
The Trinity Centre for Health Studies

Dr Emer Shelley
Department of Health and Children

Prof. Sean Strain
University of Ulster, Coleraine

SUMMARY

Cardiovascular disease, including heart disease, stroke and related diseases is the single highest cause of death in Ireland, accounting for over two in five (approximately 41%) of all deaths. High blood pressure is one of the major modifiable causal factors in the development of cardiovascular disease. In recent decades, a substantial body of evidence has emerged from observational and experimental research to suggest that high dietary salt intake is an important causal factor in the rise in blood pressure with age and in the development of essential hypertension in industrialised countries such as Ireland. There is also evidence that relatively modest reductions in salt intake have the potential to produce a significant fall in average blood pressure at a population level with a concomitant substantial impact on the burden of morbidity and mortality from cardiovascular disease.

The average daily salt intake in Ireland is high - approximately 10g in adults. Data are lacking on salt intake in children; however, data from the UK suggest that average daily salt intake in children aged 4-6 years and 7-10 years exceeds 5g and 6g, respectively. These intakes are well in excess of physiological requirements. The recommended dietary allowance (RDA) is 1.6g/70 mmol sodium (4g salt) per day for adults. Over 90% of the sodium in the diet is in the form of sodium chloride (salt); 1g of sodium is equivalent to 2.54g of salt.

It is estimated that about 15-20% of total dietary sodium intake is from discretionary sources (salt added in cooking and at table), 15% from naturally occurring sodium in unprocessed foods and about 65-70% from manufactured foods. Two food groups (meat/fish, particularly processed meats and bread) account for over 50% of salt intake from foods, with the remainder contributed by various other processed foods, including milk products, soups and sauces, spreading fats, biscuits/cakes/pastries/confectionery and breakfast cereals.

Two recent major reports on dietary salt and health: the UK Scientific Advisory Committee on Nutrition report on '*Salt and Health*' in 2003 and the US Institute of Medicine report on '*Dietary Reference Intakes for water, potassium, sodium, chloride and sulphate*' in 2004, provide a detailed and comprehensive review of the evidence on the relationship of salt and health. The Scientific Committee of the FSAI concurs with the main conclusions from these reports as follows:

- There is a direct, independent causal link between dietary salt intake and blood pressure.
- The magnitude of the effect of salt on blood pressure is significant at the clinical and population level.
- As there is a spectrum of responses to dietary salt exposures, there are no agreed criteria to identify salt sensitivity and the concept is of limited relevance.



- Children may be particularly vulnerable to the adverse effects of high salt intake. The short term benefits of salt restriction are likely to be most marked in the elderly because of the high absolute risk of hypertension-related stroke and heart disease in this age group.
- Measures taken to reduce average population salt intake will lead to lower average population blood pressure and reduce morbidity from cardiovascular disease including heart attack and stroke.
- As dietary salt is mostly derived from processed food, advice to individuals to reduce salt intake has limited impact on its own and effective measures to reduce salt intake in the population also requires a reduction in the salt content of processed foods.



RECOMMENDATIONS

General Recommendations

1. In the development of programmes for reducing salt consumption, an achievable target for the adult Irish population is a mean intake of 2.4g/100 mmol sodium (6g salt) per day. Whilst this is considered to be an achievable goal for the population at this time, it should not be regarded as an optimal or ideal level of consumption.
2. As distinct from the achievable population target, advice targeted at individual adults should reflect the RDA for sodium of 1.6g/70 mmol (4g salt) per adult per day is sufficient to meet the physiological needs of 97.5% of the population.
3. The particular vulnerability of children and the elderly to the adverse effects of high salt intake needs to be highlighted in discussion with the food industry regarding new product development and the reformulation of existing products. This should also be considered in health promotion campaigns mounted by public and private bodies.

Recommendations for the FSAI

4. The FSAI should continue to work in consultation with the food industry (manufacturers, retailers, caterers, etc) to achieve gradual, sustained and universal reductions in the salt content of processed and prepared foods. Targets should be set that are consistent with the objective of reducing the mean adult population salt intake to 6g per day within an agreed time frame.
5. The FSAI and other relevant Government bodies should support the clear and comprehensive nutritional labelling of the salt content of all processed pre-packaged food, including information on the proportion of the RDA for sodium (salt) in a single serving.
6. Consideration should be given to the mandatory labelling of foods with salt content above a specific threshold as 'high salt'. The FSAI in collaboration with relevant Government bodies should also work to ensure that the proposed EU health claims legislation sets clear guidelines for the use of claims such as 'low salt' or 'reduced salt' that are consistent with the recommendations of this report.
7. The FSAI should develop structures and methods to monitor the salt content of processed food and food prepared in the food service sector.



Recommendations for Other Agencies

8. The public sector should play a lead role in reducing population exposure to dietary salt through the development of policies and procedures to reduce the salt content of food served to staff and clients in public agencies, including hospitals and institutional care settings.
9. As part of a wider strategy for salt reduction in the diet, the Department of Health and Children and other relevant agencies should work in collaboration with consumer groups and other stakeholders to assist in the development of national information and awareness campaigns addressing:
 - the links between salt, blood pressure, heart attack and stroke
 - the salt content of processed food
 - the need to reduce the amount of salt added during cooking and at table.

Health education/promotion campaigns addressing salt intake should be integrated as appropriate with other core messages on healthy diet and lifestyle.

10. The Department of Health and Children should commission studies into accurately assessing salt intake in the general population, including relevant sub-groups such as children and the elderly.
11. The Department of Health and Children should commission regular population health monitoring studies addressing the distribution of blood pressure and the prevalence of hypertension in children and adults.

Recommendations for the Food Industry

12. The food industry should work in consultation with the FSAI and other relevant agencies to achieve a gradual, sustained and universal reduction in the salt content of processed foods and foods prepared by the food service sector.
13. The food industry should attach high priority to research aimed at addressing technological, shelf-life, preservation and taste issues in relation to the reduction of the salt content of processed food.

BACKGROUND TO THIS REPORT

In the 1999 Report of the Cardiovascular Health Strategy, *'Building Healthier Hearts'* (Department of Health and Children, 1999) it was recommended that the FSAI examine salt intake in the Irish diet and advise on national policy in this area. This recommendation reflects the current international scientific consensus that dietary salt intake is an important factor in the development of hypertension, a major determinant of risk of cardiovascular disease, including myocardial infarction (heart attack) and stroke. This report addresses the following issues:

1. The importance of hypertension as a cause of morbidity and mortality in Ireland.
2. The scientific evidence linking high dietary salt consumption to increased blood pressure.
3. The role of salt relative to other factors in the development of hypertension
4. Intake and sources of sodium (salt) in the Irish diet.
5. Recommended levels of salt intake.
6. Recommendations for measures to achieve target reductions in salt intake in children and adults.

The Committee has undertaken a detailed review of the national and international literature on salt intake and health, with particular reference to the link between salt and blood pressure. Two recent major reports on dietary salt and health provided detailed and comprehensive reviews of current research in this area: the UK Scientific Advisory Committee on Nutrition (SACN) report on *'Salt and Health'* (SACN, 2003) and the US Institute of Medicine (IOM) report on *'Dietary Reference Intakes for water, potassium, sodium, chloride and sulphate'* (US IOM, 2004). Given the recent publication of these reports, the Committee agreed that a further in-depth review of the scientific evidence is not required at this time. Details of these reports are not reproduced here; however, references to websites where the summaries of each of these reports can be accessed are contained in the list of references. This report will give a brief overview of the scientific evidence linking high dietary salt consumption with increased blood pressure.

CHAPTER I. HYPERTENSION AND CARDIOVASCULAR DISEASE IN IRELAND

Definition of Hypertension

Blood pressure is a continuous, normally distributed variable in the population. Hypertension is therefore a quantitative phenomenon in which the definition of abnormality is arbitrary and pragmatic. In essence, we define hypertension as the blood pressure level at which the benefits of intervention are adjudged to outweigh the costs. The current internationally agreed cut-off point for normal blood pressure is at a systolic blood pressure (SBP) of less than 140 mmHg and diastolic blood pressure (DBP) of less than 90 mmHg. An additional category of 'high normal' blood pressure (SBP between 120 and 139 or DBP between 85 and 89 mmHg) is widely accepted on the basis that persons with blood pressures within these ranges are at increased risk of future 'hypertension' and cardiovascular events relative to those with 'normal' blood pressure, and are likely to benefit from non-pharmacological interventions (Chobanian, 2003).

Prevalence of Hypertension

It is estimated that 24% of the US adult population, meet current internationally agreed diagnostic criteria for hypertension, with substantially higher prevalence rates, approaching 70%, in persons aged 70 years and older (Burt *et al*, 1995). In the 2002 Health Survey for England, mean SBP for men was 135 mmHg and for women 131 mmHg (Health Survey for England, 2002). In this national survey, mean SBP increased with age in both men and women, rising from 128 mmHg in men aged 16-24 to 146 mmHg in men aged 75 and over and

from 120 mmHg to 152 mmHg in women. In England, it is estimated that 37% of men and 34% of women have hypertension (defined here as a SBP of 140 mmHg or over, or a DBP of 90 mmHg or over) or are being treated for hypertension. The prevalence of hypertension increases with age in both sexes; it is estimated that less than one in twenty women aged 16-24 are hypertensive, compared to more than half aged 55-64 and just under three-quarters aged 65-74. Just under 80% of men and 70% of women with hypertension are not receiving treatment. Of those that are treated, over 60% remain hypertensives (Health Survey for England, 2002).

By contrast with the situation in the UK, there are limited contemporary data on the distribution of blood pressure and the prevalence of hypertension in the Irish population. On the basis of the recent data from England we can estimate that in Ireland, at least 50% of the population aged 50 years and older are hypertensive (Health Survey for England, 2002). In a cross-sectional study carried out in 1998, involving 1,018 men and women aged 50 to 69 years, drawn from General Practice Registers in Cork and Kerry, Creagh and colleagues reported a prevalence of hypertension of 50% in males and 43.5% in females. Only 38% of those meeting current international criteria for hypertension had been previously diagnosed and were on treatment (Creagh *et al*, 2002).

Hypertension and Cardiovascular Disease

Heart disease and stroke are the first and third leading causes of death in most developed countries, including Ireland, and it is estimated that they will rank as first and fourth respectively as causes of the global burden of disease by the year 2020 (Lopez & Murray, 1998). Hypertension is one of the most important modifiable risk factors for both coronary heart disease and stroke. Other major factors include smoking, diet, physical inactivity and diabetes. Data from large-scale observational studies involving follow-up of hundreds of thousands of individuals in the general population show a consistent, linear relation between usual blood pressure levels and risk of coronary heart disease (CHD) and stroke (MacMahon *et al*, 1990; Lewington *et al*, 2002). Elevated blood pressure is also a major predictor of heart failure and of end-stage renal failure (Whelton *et al*, 1996). The findings from randomised controlled trials of blood pressure lowering drugs also provide clear evidence that blood pressure levels are directly and continuously related to the risks of stroke, CHD and other adverse outcomes across the entire distribution of blood pressure in the population (Collins, 1990). Because of the continuous relation between blood pressure and risk, small downward shifts in mean population blood pressure will produce substantial reductions in both the proportion of hypertensives in the upper tail of the distribution and in hypertension related morbidity and mortality (Stamler *et al*, 1989). This has profound implications for public health policy in relation to salt and blood pressure.

CHAPTER 2. OVERVIEW OF EVIDENCE ON SALT AND BLOOD PRESSURE

The contribution of dietary factors such as salt to the rise in blood pressure with age and the development of essential hypertension have been difficult to elucidate because of the poor precision (or reliability) with which dietary exposures are measured in free-living subjects and the limited range of dietary exposures in most populations. There is also the problem of multicollinearity (i.e. the tendency for dietary elements such as sodium, potassium and fat to be intercorrelated at both the individual and population level) (Perry *et al*, 1994). Despite these difficulties, however, the evidence that salt intake plays a critical role in blood pressure regulation is now overwhelming (MacGregor and de Wardener, 2002; Elliott and Stamler, 2002).

The evidence comes from diverse sources, including:

- observational epidemiologic studies (ecological, cross-sectional and migration studies)
- animal and genetic models
- a trial in infants with long term follow-up
- a community intervention study carried out in Portugal
- intervention studies in normotensives (people with normal blood pressure) and hypertensives (people with high blood pressure) and in studies involving middle-aged and elderly population samples.

In an international study including over 10,000 subjects (men and women aged 20–59 years) in 52 different population groups in 32 countries, positive associations between urinary sodium excretion (a marker of salt intake) and blood pressure were observed within and between populations (INTERSALT, 1988; Elliott *et al*, 1996). Within populations, those with higher sodium excretion tended to have higher blood pressure. In the across population analysis (i.e. ecological analysis comparing blood pressure levels in 32 different countries) populations with higher mean sodium excretion had higher mean blood pressures. In men and women at all ages it was estimated that a 100 mmol/day increase in sodium intake was associated with an average increase in systolic blood pressure of up to 6 mmHg. Estimates of association were larger for older people (aged 40–59 years) than for younger people (aged 20–39 years). One of the key findings from this study was a consistent and highly significant association of sodium excretion across populations with the slope or rise of blood pressure with age (INTERSALT, 1988; Elliott *et al*, 1996).

In virtually all mammals, high blood pressure is caused or aggravated by a high salt intake, e.g. it has been shown in chimpanzees that an increase in salt intake from 0.5g daily (their usual intake) to a level between 9g and 15g daily (our usual intake), leads to substantial and reversible increases in blood pressure (Denton, 1995).

In a randomised controlled trial that involved 500 newborn infants in the Netherlands, it was found that infants who were given formula milk and solids with reduced salt content had significantly lower blood pressure at six months of age relative to a control group on a standard infant diet (Hofman, 1983). Intriguingly, on re-examination of a sub-group of these children at age 15 years, there was evidence that the beneficial effects of early salt restriction on blood pressure persisted into adolescence (Geleijnse, 1996). The data from this study combined with the findings from INTERSALT on the positive association between average population salt intake and the rise of blood pressure with age suggest that children are especially vulnerable to adverse effects from high salt intake (INTERSALT, 1988; Elliott *et al*, 1996).

In the community intervention trial in Portugal, the salt intake of an entire village was reduced by reducing salt in cooking and in processed food, including bread. At the end of the observation period, blood pressure was significantly lower than in a control village (Forte *et al*, 1989).

Meta-analyses of Salt Restriction Intervention Studies

The best evidence for the role of salt in blood pressure comes from randomised controlled trials. Feng and McGregor have recently reported the findings from a meta-analysis, based on 17 trials of modest reductions in salt intake in hypertensives and 11 similar trials in normotensives (Feng and McGregor, 2002). Trials of very short duration (one week or less) and acute salt loading and salt depletion experiments were excluded from the meta-analyses. The findings from this meta-analysis are consistent with those of Law and colleagues (1991). The median reduction in salt intake in the trials was approximately 5g per day and this was associated with an average fall in blood pressure of 4.96/2.73 mmHg in hypertensives and 2.03/0.97 mmHg in normotensives. In linear regression analyses, a reduction of 100mmol per day (6g) in salt intake predicted a fall in blood pressure of 7.1/3.9 mmHg in hypertensives and 3.6/1.6 mmHg in normotensives (Feng and McGregor, 2002). It is estimated that an average reduction in blood pressure of this magnitude in the general population of most Western countries would reduce the incidence of stroke by 24% and the incidence of ischemic heart disease by 18% (Stamler *et al*, 1989; Cook *et al*, 1995). This would lead to a reduction in stroke deaths of approximately 600 per year in Ireland (with a similar reduction in the number of non-fatal disabling strokes) and a reduction in deaths from ischaemic heart disease of approximately 1,100 per year. These effects are substantial in population health terms and are consistent with the hypothesis that even a modest reduction in average dietary salt intake at the population level is likely to produce substantial falls in stroke and coronary heart disease mortality.

At least six meta-analyses of salt restriction trials have been conducted to date (summarised in the SACN 2003 report), a number of which reported less marked effects of salt restriction on blood pressure. It is noteworthy, however, that in the dissenting meta-analyses, significant effects of salt restriction on blood pressure were detected in both hypertensives and normotensives but the effect in normotensives was judged to be of insufficient magnitude to be of public health importance (Midgley *et al*, 1996; Cutler *et al*, 1997; Graudal *et al*, 1998).

Salt Restriction in the Elderly

Reductions in salt intake may be of particular benefit in the elderly. In a randomised controlled trial involving men and women aged 60–78 years, a reduction in daily salt intake from 10 to 5g for a period of one month was associated with an average fall in SBP of 7 mmHg. These effects, which were seen in normotensive and hypertensive subjects, translate into an estimated 36% reduction in stroke risk over a five year period in this age group (Cappuccio *et al*, 1997). Given the high underlying incidence of stroke in the elderly, a reduction in stroke incidence of this magnitude (more than one third) would represent a public health triumph. In this, as in other similar studies, there was no evidence of a distinct sub-group of so-called salt-sensitive subjects. This concept that a minority of the population may be salt-sensitive, with the rest of the population being relatively unaffected, has now been discredited (US IOM, 2003).

The Dash Sodium Trial

Observational epidemiological studies and meta-analyses based on trials of varying methodological rigour in different populations and patient groups have provided fertile ground for controversy on the link between salt and blood pressure. In this context, the findings from the DASH Sodium study (dietary approaches to stop hypertension) are particularly important (Sacks *et al*, 2001). This was a well conducted, three month intervention trial, too brief to examine morbidity, but long enough to detect clear effects of diet on blood pressure. Four hundred and twelve participants, ranging in age from 37 - 59 years, were randomly assigned to a control diet typical of the U.S usual diet or the DASH diet (rich in fruit, vegetables, low-fat dairy products, with a reduced total and saturated fat intake (Appel *et al*, 1997). The study included three levels of sodium intake:

- high (150 mmol sodium/9g salt)
- intermediate (100 mmol sodium/6g salt)
- low (50 mmol sodium/3g salt).

The results of the DASH sodium study showed a graded linear relation between salt intake and blood pressure from high through intermediate to low intakes. The effects were seen in patients on both the control and DASH diets, in those with and without hypertension, in both sexes and across racial groups. Among normotensives on the control diet, lower salt intake (50 mmol sodium per day) versus higher (150 mmol sodium per day) decreased blood pressure by 7.0/3.8 mmHg in those older than 45 years of age and by 3.7/1.5 mmHg in those 45 years of age or younger. Of particular importance in this study was the finding that the combination of salt restriction with the DASH study diet produced additive effects on blood pressure in both hypertensives and normotensives. When compared with the control diet with a high sodium level, the DASH diet with a low sodium level led to a mean SBP that was 7.1 mmHg lower in participants without hypertension and 11.5 mmHg lower in participants with hypertension (Sacks *et al*, 2001).

Salt, Left Ventricular Hypertrophy (Cardiac Enlargement) and Stroke

It is suggested that associations between sodium intake and blood pressure in observational studies are attenuated by measurement imprecision reflecting large intra-subject variation in salt intake and blood pressure. Therefore, stronger associations between sodium intake and stable indices of hypertensive end-organ damage such as left ventricular hypertrophy (enlargement of the heart) could be expected. Consistent with this hypothesis, there is accumulating evidence that high salt intake predicts left ventricular hypertrophy (Schneider *et al*, 1988). This association persists in analyses adjusted for blood pressure. There is also accumulating evidence that in overweight persons higher dietary salt intake is associated with increased risk of CHD, stroke and overall mortality (Tuomilehto *et al*, 2001; He *et al*, 1999). This evidence is summarised in Annex 1 of the SACN report.

CHAPTER 3. THE ROLE OF SALT RELATIVE TO OTHER FACTORS IN THE DEVELOPMENT OF HYPERTENSION

Causes of Hypertension

The fundamental problem in hypertension is the tendency for blood pressure to rise with age. The extent of the blood pressure rise with age and thus the occurrence of hypertension varies considerably both within populations and between populations world-wide (Perry, 1994). Thus hypertension is as much a disorder of populations as of individuals (Rose, 2001).

The development of hypertension reflects a complex and dynamic interaction between genetic and environmental causal factors. Differences in genetic susceptibility probably account for much of the blood pressure variation within a population such as Ireland where we are all exposed to a broadly similar environment, whereas differences in environmental factors largely determine variation in blood pressure levels between populations and within populations over time. The findings from the DASH trial highlight the importance of dietary factors in the development of hypertension, with significant beneficial effects of fruit and vegetables (important sources of potassium, magnesium and fibre) and low-fat dairy products (important sources of calcium and potassium) on blood pressure in addition to the effects of reducing salt intake. The other major determinants of blood pressure levels in the population are overweight and obesity, physical inactivity and high alcohol consumption (Perry *et al.* 1994). As these factors are highly inter-correlated, the relative contribution of each to the overall burden of hypertension cannot be precisely estimated.

Salt is a high priority issue for Ireland in the development of public policy initiatives to control blood pressure for several reasons:

current trends in other causal factors (notably obesity and physical inactivity) are unfavourable and are unlikely to be reversed in the short term, we have high mortality and morbidity rates from heart disease and stroke relative to other developed countries and, working in collaboration with the food industry, policy options are available to us which will effectively reduce population exposure to dietary salt without major changes in consumer behaviour. It should also be noted that control of dietary salt intakes at the population level offers an extremely cost effective approach to the control of hypertension relative to alternative, clinically oriented approaches to the problem: disease labelling, poly-pharmacy and clinical supervision over several decades for approximately 50% of the population. In the WHO Global burden of disease study it has been estimated that societal level action to stimulate a reduction in salt content of processed foods could avert over 21 million disability adjusted life years (DALYS) per year worldwide (Murray *et al.*, 2003).

Intake and Food Sources of Sodium/Salt in Ireland

Sodium is present naturally in foods and drinking water. As salt (sodium chloride), it is added to foods during processing, cooking and at the table. The main reasons for addition of salt in food manufacture are for flavour, texture and preservation. Sodium chloride (salt) contains about 40% sodium by weight (1g sodium chloride = 0.4g sodium). Other sodium salts are added to foods during manufacture, e.g. sodium nitrate to cured meats for preservation and sodium bicarbonate to bread for texture, but generally in much lower amounts than sodium chloride.

Major dietary sources of sodium are from salt added during the processing and manufacture of foods (non-discretionary) and salt added to food during cooking and at the table (discretionary). It is estimated that about 15-20% of total dietary sodium intake is from discretionary sources, 15% from naturally occurring sodium in unprocessed foods and about 65-70% from manufactured foods (British Nutrition Foundation, 1994; Bull & Buss 1990; Flynn *et al*, 1990; SACN, 2003).

Average daily sodium (salt) intake from foods in Irish adults has been estimated as 3.25g (8.3g salt) (Table 1), with a 95th percentile of 5.23g (13.3g salt). This estimate does not allow for all additions in cooking or any additions at table – assuming about 15-20% of total dietary sodium intake is from discretionary sources would give a total daily sodium intake of about 4g (10g salt). This is similar to the estimate of total daily sodium intake of 3.8g (about 9.5g salt) obtained for UK adults using the 24-hour urinary excretion method, which is considered a more accurate method to determine total salt intake (Henderson *et al*, 2003).

In a study carried out in University College, Cork, in a sample of 114 adult hospital outpatients (77 males and 37 females) with Type 2 diabetes, mean daily sodium intake, estimated from 24-hour urinary output, for all patients was 166 mmol (about 10g salt) (MacLeod *et al*, 2005). Intakes were significantly lower in females, 144 mmol (about 8g salt) compared to males 177.2 mmol (about 10g salt). Dietary assessment using 24-hour recalls indicated that the main food sources of salt were breads and cereals (38%) and meat, including processed meats (35%), with less than 10% from salt added at the table. This study suggests that patients with Type 2 diabetes have a high salt intake, despite clear motivating factors and dietary counselling.

There are no data on salt intakes of children and adolescents in Ireland. In the UK, average daily sodium (salt) intake from foods in 4 to 18 year olds has been estimated as: 4-6 yr, 1.97g (5.0g salt); 7-10 yr, 2.28g (5.8g salt); 11-14 yr, 2.49g (6.31g salt); and 15-18 yr, 2.79g (7.1g salt) (Gregory *et al*, 2000). Again, this estimate does not allow for additions in cooking and at table.

The main food sources of sodium for adults in Ireland are listed in Table 1. Meat, fish and their products, particularly processed meats, provide almost 30% of total sodium (salt) intake, with a further 26% provided by bread and rolls. Other contributing foods include: milk and milk products, soups, sauces and miscellaneous foods, spreading fats, biscuits/cakes/pastries/confectionery and breakfast cereals.

Table 1. Mean daily Sodium Intake from Foods in Irish Adults Aged 18-64 Years by Food Group*

| Food/Food Category | g/day | % total |
|---|--------------|----------------|
| Meat & fish | 0.97 | 29.8 |
| <i>Cured/processed meats</i> | <i>0.67</i> | <i>20.5</i> |
| <i>Meat/meat dishes</i> | <i>0.23</i> | <i>6.9</i> |
| <i>Fish/fish dishes</i> | <i>0.08</i> | <i>2.4</i> |
| Bread & rolls | 0.84 | 25.9 |
| Milk & milk products | 0.27 | 8.5 |
| <i>Cheese</i> | <i>0.12</i> | <i>3.5</i> |
| Soups, sauces & miscellaneous foods | 0.23 | 7.0 |
| Spreading fats | 0.19 | 5.9 |
| Biscuits/cakes/pastries/confectionary | 0.15 | 4.5 |
| Breakfast cereals | 0.14 | 4.2 |
| <i>Ready-to-eat breakfast cereals</i> | <i>0.13</i> | <i>4.1</i> |
| <i>Other</i> | <i>trace</i> | <i>0.1</i> |
| Vegetables/processed vegetables | 0.13 | 4.0 |
| <i>Processed vegetables/vegetable dishes</i> | <i>0.04</i> | <i>1.1</i> |
| Savouries (e.g. pizza, mixed pasta dishes) | 0.095 | 2.9 |
| Egg/egg dishes | 0.049 | 1.5 |
| Desserts | 0.035 | 1.1 |
| Other foods | 0.15 | 4.7 |
| Total | 3.25 | 100.0 |

* Estimated from the North/South Ireland Food Consumption Survey (IUNA, 2001) for Republic of Ireland only (n=776), excluding under-reporters by the method of Black (2000)

Salt Taste Thresholds

There is considerable evidence that salt taste thresholds fall with decreasing intake. Food with a high salt content becomes unpalatable within 4-6 weeks of adopting a low salt diet. The Scientific Committee is of the view that this concept of adaptation to reduced salt intake should be communicated clearly to consumers. There is also evidence that significant incremental reductions in the salt concentration of processed food (up to 10% per year) can be achieved without adverse effects on taste (MacGregor & Sever, 1996). This has clear and significant implications for the food industry.

CHAPTER 4. RECOMMENDED LEVELS OF SODIUM INTAKE

In 1991, the UK Committee on the Medical Aspects of Food and Nutrition Policy (COMA) set a reference nutrient intake (the term RDA, recommended dietary allowance is used in this document) for sodium of 1.6g/70 mmol (4g salt), per day to meet the needs of 97.5% of the population. This was endorsed by the UK Scientific Advisory Committee on Nutrition (SACN, 2003), which succeeded COMA in 2003. The Food Standards Agency (FSA) in the UK and the Institute of Medicine (IOM) National Academy of Science in the USA recommend an upper level of no higher than 2.4g/100 mmol sodium (6g salt) per day. This is considered a population target not an optimal or ideal level of consumption.

The Scientific Committee endorses these recommendations, emphasising that the RDA of 1.6g sodium (4g salt) per day should form the basis of advice targeted at individuals as distinct from the population health target of a mean salt intake of 6g per day. The latter target is regarded as an achievable (as distinct from ideal) goal for the population at this time. It represents a substantial reduction in salt intakes and will require continued co-operation of food manufacturers, caterers and retailers as well as increased awareness of individuals of the need to reduce their own salt intake. This target should be reviewed as average salt intake falls in the population, thereby reducing salt taste thresholds.

Salt intake in children needs to be proportionally lower than in adults (based on body weight). The Committee also endorses the target levels for average daily salt intake recommended for children in the UK SACN report on 'Salt and Health' (2003):

| | |
|-----------------|-----|
| Age 0-6 months | <1g |
| Age 7-12 months | 1g |
| Age 1-3 years | 2g |
| Age 4-6 years | 3g |
| Age 7-10 years | 5g |
| Age 11-14 years | 6g |

Potential Interventions to Reduce Population Salt Intake

The Committee discussed a wide range of potential initiatives that might be considered in developing measures to reduce population salt intake. Among potential initiatives discussed were the following:

- Advice to individuals and public education programmes to promote a reduction of salt added table in cooking and at table.
- Voluntary reduction of salt in processed food by industry.
- Voluntary reduction of salt in food prepared by the catering industry.

- 
- Promotion of the use of low salt substitutes and herbs/spices.
 - Regulation of nutritional labelling of the salt content of all processed food.
 - Regulation of the salt content in processed food.

Worldwide, interventions have ranged from non-specific advice to the public to reduce dietary salt intake to voluntary collaborations with the food and catering industries to reduce salt content in processed and pre-prepared foods. It is recognised that achievement of the population average salt intake targets will require substantial effort and all of the above approaches should be explored.

However, regarding the use of low sodium salt, the Committee was of the opinion that this approach to reducing population dietary sodium cannot be endorsed at this time. Concerns were raised about the possible vulnerability of certain population sub-groups (including those with Type 1 diabetes, chronic renal insufficiency, end stage renal disease, severe heart failure and adrenal insufficiency) to high potassium load from these salt substitutes. It was also noted that the use of salt substitutes does not address the need to reduce salt taste thresholds in the population.

A systematic review by Hooper *et al* (2002) assessed the long-term effects of advice to reduce dietary salt in adults with and without hypertension. Eleven trials were included with follow-up from 6 months to 7 years. The significant blood pressure reductions observed at 6-12 months, were not sustained over time. As most dietary salt intake originates from processed food, this illustrates the difficulties faced by individuals in maintaining a low salt diet and supports the view that a targeted individual approach to salt reduction will have limited impact without concomitant reduction in salt content of foods. A population-based approach is required to achieve a sustained reduction in salt intake.

CHAPTER 5. CONCLUSIONS

Salt intake in the Irish population is high. There is convincing evidence of a causal link between high salt intake and hypertension and of the benefit of reducing salt intake at population level. Hypertension is a significant cause of illness and death in Ireland. With the co-operation of the food industry and increased awareness of individuals of the need to reduce their own salt intake, it is possible to reduce adult population salt intake from an average of about 10 grams per day to about 6 grams per day. There is now a scientific consensus that a reduction of this magnitude will lead to a significant fall in blood pressure and substantial reductions in suffering and death from heart attack, stroke and related conditions.

CHAPTER 6. MAJOR RECOMMENDATIONS

General Recommendations

1. In the development of programmes for reducing salt consumption, an achievable target for the adult Irish population is a mean intake of 2.4g/100 mmol sodium (6g salt) per day. Whilst this is considered to be an achievable goal for the population at this time, it should not be regarded as an optimal or ideal level of consumption.
2. As distinct from the achievable population target, advice targeted at individual adults should reflect the RDA for sodium of 1.6g/70mmol (4g salt) per adult per day is sufficient to meet the needs of 97.5% of the population.
3. The particular vulnerability of children and the elderly to the adverse effects of high salt intake needs to be highlighted in discussion with the food industry regarding new product development and the reformulation of existing products. This should also be considered in health promotion campaigns mounted by public and private bodies.

Recommendations for the FSAI

4. The FSAI should continue to work in consultation with the food industry (manufacturers, retailers, caterers, etc) to achieve gradual, sustained and universal reductions in the salt content of processed and prepared foods. Targets should be set that are consistent with the objective of reducing the mean adult population salt intake to 6g per day within an agreed time frame.
5. The FSAI and other relevant Government bodies should support the clear and comprehensive nutritional labelling of the salt content of all processed pre-packaged food, including information on the proportion of the RDA for sodium (salt) in a single serving.
6. Consideration should be given to the mandatory labelling of foods with salt content above a specific threshold as 'high salt'. The FSAI in collaboration with relevant Government bodies should also work to ensure that the proposed EU health claims legislation sets clear guidelines for the use of claims such as 'low salt' or 'reduced salt' that are consistent with the recommendations of this report.
7. The FSAI should develop structures and methods to monitor the salt content of processed food and food prepared in the food service sector.

Recommendations for Other Agencies

8. The public sector should play a lead role in reducing population exposure to dietary salt through the development of policies and procedures to reduce the salt content of food served to staff and clients in public agencies, including hospitals.
9. As part of a wider strategy for salt reduction in the diet, the Department of Health and Children and other relevant agencies should work in collaboration with consumer groups and other stakeholders to assist in the development of national information and awareness campaigns addressing:
 - the links between salt, blood pressure, heart attack and stroke
 - the salt content of processed food
 - the need to reduce the amount of salt added during cooking and at table.

Health education/promotion campaigns addressing salt intake should be integrated as appropriate with other core messages on healthy diet and lifestyle.

10. The Department of Health and Children should commission studies into salt intake in the general population, including relevant sub-groups such as children and the elderly.
11. The Department of Health and Children should commission regular population health monitoring studies addressing the distribution of blood pressure and the prevalence of hypertension in children and adults.

Recommendations for the Food Industry

12. The food industry should work in consultation with the FSAI and other relevant agencies to achieve a gradual, sustained and universal reduction in the salt content of processed foods and foods prepared by the food service sector.
13. The food industry should attach high priority to research aimed at addressing technological, shelf-life, preservation and taste issues in relation to the reduction of the salt content of processed food.

REFERENCES

- Appel, L.J., Moore, T.J., Obarzanek, E., Vollmer, W.M., Svetkey, L.P., Sacks, F.M., Bray, G.A., Vogt, T.M., Cutler, J.A., Windhauser, M.M., Lin, P.H., Karanja, N.** (1997) A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group 1997. *N Engl J Med* 336:1117-1124.
- Black, A.E.** (2000) Critical evaluation of energy intake using the Goldberg cut-off for energy intake: basal metabolic rate. A practical guide to its calculation, use and limitations. *International Journal of Obesity* 24(9):1119-1130.
- British Nutrition Foundation** (1994) Salt in the Diet. Briefing Paper.
- Bull, N.L., Buss, D.H.** (1990) Contribution of foods to sodium intakes. *Proc. Nutr. Soc* 39: 40A.
- Burt, V.L. C.J., Higgins, M., Horan, M.J., Labarthe, D., Whelton, P., Brown, C., Rocella, E.J.** (1995) Trends in prevalence, awareness, treatment and control of hypertension in the adult US population. Data from the Health Examination Surveys 1960 to 1991. *Hypertension* 26:60-69.
- Chobanian, A.V., Bakris, G.L., Black, H.R., Cushman, W.C., Green, L.A., Izzo, J.L. Jr. et al.** (2003) National High Blood Pressure Education Program Coordinating Committee. Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension* 42:1206-1252.
- Cappuccio, F.P., Markandu, N.D., Carney, C., Sagnella, G.A., MacGregor, G.A.** (1997) Double blind randomised trial of modest salt reduction in older people. *Lancet* 350: 850-854.
- Collins, R., Peto, R., MacMahon, S., Hebert, P., Fiebach, N.H., Eberlein, K.A., et al.** (1990) Blood pressure, stroke, and coronary heart disease. Part 2, Short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. *Lancet* 335:827-38.
- Cook, NR., Cohen, J., Hebert, PR., Taylor, J.O., Hennekens, C.H.** (1995) Implications of small reductions in diastolic blood pressure for primary prevention. *Arch Intern Med* 155:701-709.
- Creagh, D., Neilson, S., Collins, A., Colwell, N., Hinchion, R., O'Halloran, D., Perry, I.J.** (2002) Established cardiovascular disease and CVD risk factors in a primary care population of middle-aged Irish men and women. *Ir Med J* 95:298-301.
- Cutler, J.A., Follmann, D., Allender, P.S.,** (1997) Randomised trials for sodium reduction: an overview. *Am J Clin Nutr* 65 (2 Suppl):643S-651S.
- Denton, D., Weisinger, R., Mundy, N.I. et al.** (1995) The effect of increased salt intake on blood pressure of chimpanzees. *Nature Medicine* 1:1009-16.

Department of Health and Children
(1999) Building Healthier Hearts
http://www.dohc.ie/publications/building_healthier_hearts.html

Elliott, P., Stamler, J., Nichols, R., Dyer, A.R., Stamler, R., Kesteloot, H., Marmot, M. (1996) Intersalt revisited: further analyses of 24 hour sodium excretion and blood pressure within and across populations. Intersalt Cooperative Research Group. *BMJ* 312:1249-1253.

Elliott, P., Stamler, J. (2002) Evidence on salt and blood pressure is consistent and persuasive. *Int J Epidemiol* 31:316-319.

Feng, J.H., MacGregor, G.A. (2002) Effect of modest salt reduction on blood pressure. A meta-analysis of randomised trials: implications for Public Health. *J Hum Hypertens* 16:761-770.

Flynn, A., Shortt, C., Morrissey, P.A. (1990) Sodium and potassium intakes in Ireland. *Proc Nutr Soc* 49:323-32.

Forte, J.G., Miguel, J.M., Miguel, M.J., de Padua, F., Rose, G. (1989) Salt and blood pressure: a community trial. *J Hum Hypertens* 3:179-84.

Geleijnse, J.M., Hofman, A., Witteman, J.C.M., Hazebroek, A.A.J.M., Valkenburg, H.A., Grobbee, D.E. (1996) Long-term effects of neonatal sodium restriction on blood pressure. *Hypertension* 29:913-917.

Graudal, N.A., Galloe, A.M., Garred, P. (1998) Effects of sodium restriction on blood pressure, renin, aldosterone, catecholamines, cholesterol, and triglyceride. A meta-analysis. *J Am Med Assoc* 279:1383-1391.

Gregory, J., Lowe, S., Bates, C.J., Prentice, A., Jackson, L.V., Smithers, G., Wenlock, R., Farron, M. (2000) National Diet and Nutrition Survey: young people aged 4 to 18 years. *Volume 1: Report of the diet and nutrition survey*. TSO, London.

He, J., Ogden, L.G., Vupputuri, S., Bazzano, L., Loria, C., Whelton, P.K. (1999) Dietary sodium intake and subsequent risk of cardiovascular disease in overweight adults. *JAMA* 282:2027-2034.

Health Survey for England, Joint Health Surveys Unit (2002) National Centre for Social Research, Department of Epidemiology and Public Health at the Royal Free and University College Medical School. <http://www.official-documents.co.uk/document/deps/doh/survey02/hse02.htm>

Henderson, L., Gregory, J., Irving, K and Swan, G. (2002) The National Diet and Nutrition survey: adults aged 19-64 years. *Volume 2. Energy, protein, carbohydrate, fat and alcohol intake*. London:TSO.

Hofman, A., Hazelbroek, A., Valkenburg, H.A. (1983) A randomised trial of sodium intake and blood pressure in newborn infants. *JAMA* 250:370-3.



Hooper, L., Bartlett, C., Smith, G.D., Ebrahim, S. (2002) Systematic review of long term effects of advice to reduce dietary salt in adults. *BMJ* 325:628-637.

INTERSALT Cooperative Research Group. (1988) Intersalt: an international study of electrolyte excretion and blood pressure. Results for 24 hour urinary sodium and potassium excretion. *BMJ* 297:319-328.

Food Safety Promotion Board (2001) IUNA (Irish Universities Nutrition Alliance) North/South Ireland food consumption survey. Report. Dublin. www.iuna.net.

Law, M.R., Frost, C.D., Wald, N.J. (1991) By how much does dietary salt reduction lower blood pressure? III--Analysis of data from trials of salt reduction. *BMJ* 302:819-24.

Lewington, S., Clarke, R., Qizilbash, N., Peto, R., Collins, R. (2002) Age-specific relevance of usual blood pressure to vascular mortality: A Meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 360:1903-1913.

Lopez, A.D., Murray, C. C. J. L. (1998) The global burden of disease, 1990-2020. *Nature Medicine* 4:1241-1243.

MacGregor, G., de Wardener, H.E. (2002) Salt, blood pressure and health. *Int J Epidemiol* 31:320-327.

MacGregor, G.A., Sever, P.S. (1996) Salt--overwhelming evidence but still no action: can a consensus be reached with the food industry? *BMJ* 312:1287-1289.

MacLeod, F., Collins, M., Hinchion, R., Perry, I.J. (2005) Salt intake in Irish patients with Type 2 Diabetes (*In preparation*).

MacMahon, S., Peto, R., Culter, J., Collins, R., Sorlie, P., Neaton, J., et al. (1990) Blood pressure, stroke and coronary heart disease. Part I, prolonged differences in blood pressure: prospective observational studies corrected for the regressions dilution bias. *Lancet* 335:765-774.

Midgeley, J.P., Matthew, A.G., Greenwood, C.M., Logan, A.G. (1996) Effect of reduced dietary sodium on blood pressure: a meta-analysis of randomized controlled trials. *JAMA* 275 1590-1597.

Murray, C.J.L., Lauer, J.A., Hutubessy, R.C.W., Niessen, L., Tomijima, N., et al. (2003) Effectiveness and costs of interventions to reduce systolic blood pressure and cholesterol: A global and regional analysis on reduction of cardiovascular-disease risk. *Lancet* 361:717-725.

Perry, I.J., Whinup, P.H., Shaper, A.G. (1994) Environmental factors in the development of essential hypertension. *Br Med Bull* 50:246-59.

Rose, G. (2001) Sick individuals and sick populations. *Int J Epidemiol* 30:427-432 (Reiteration of *Int J Epidemiol* 1985;14:32–38).

Sacks, F.M., Svetkey, L.P., Vollmer, W.M., Appel, L.J., Bray, G.A., Harsha, D., et al. (2001) Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *N Engl J Med* 344:3-10

Schneider, R.E., Messerli, F.H., Garavaglia, G.E., Nunez, B.D. (1988) Dietary salt intake: A determinant of cardiac involvement in essential hypertension. *Circulation* 78:1951-1956.

Scientific Advisory Committee on Nutrition (2003) Salt and Health Report <http://www.food.gov.uk/multimedia/pdfs/sacnforumreport.pdf>

Stamler, J., Rose, G., Stamler, R., Elliott, P., Dyer, A., Marmot, M. (1989) INTERSALT study findings. Public health and medical care implications. *Hypertension* 14:570-7.

Tuomilehto, J., Jousilahti, P., Rastenyte, D., Moltchanov, V., Tanskanen, A., Pietinen, P., Nissinen, A. (2001) Urinary sodium excretion and cardiovascular mortality in Finland: a prospective study. *Lancet* 357:848-851.

United States Institute of Medicine. (2004) Dietary reference intakes for water, potassium, sodium, chloride and sulphate. Panel on dietary reference intakes for electrolytes and water, Standing committee on the scientific evaluation of dietary reference intakes. http://www.nap.edu/execsumm_pdf/10925.pdf

Whelton, P.K., Perneger, T.V., He, J., Klag, M.J. (1996) The role of blood pressure as a risk factor for renal disease: A review of the epidemiological evidence. *J Hum Hypertens* 10:683-689



NOTES



NOTES



Food Safety Authority of Ireland Údarás Sábháilteachta Bia na hÉireann
Abbey Court, Lower Abbey Street, Cúirt na Mainistreach, Sráid na Mainistreach Íocht.,
Dublin 1 Baile Átha Cliath 1

Telephone: +353 1 817 1300
Facsimile: +353 1 817 1301
E-mail: info@fsai.ie
Website: www.fsai.ie

€15.00